

Referenties - Wetenschappelijke onderbouwing hulpmiddelen

VierDimensionale Klachtenlijst (4DKL)

- de Croon e.a. (2005). Drie vragenlijsten voor diagnostiek van depressie en angststoornissen. *Tijdschr Bedrijfs Verzekeringsgeneesk*;13(4):98-103.
- Langerak (2012). A validation study of the Four-Dimensional Symptom Questionnaire (4DSQ) in insurance medicine. *Work*;43(3):369-80.
- Terluin e.a. (2006). The Four-Dimensional Symptom Questionnaire (4DSQ): a validation study of a multidimensional self-report questionnaire to assess distress, depression, anxiety and somatization. *BMC Psychiatry*;22;6:34.
- Terluin e.a. (2009). Detecting depressive and anxiety disorders in distressed patients in primary care; comparative diagnostic accuracy of the Four-Dimensional Symptom Questionnaire (4DSQ) and the Hospital Anxiety and Depression Scale (HADS). *BMC Fam Pract*;23;10:58.
- Terluin e.a. (2016). The Four-Dimensional Symptom Questionnaire (4DSQ) in the general population: scale structure, reliability, measurement invariance and normative data: a cross-sectional survey. *Health Qual Life Outcomes*;15;14(1):130.

Anamneselijst incontinentie

- Anamneselijst urine-incontinentie. Zorg voor beter (jaar onbekend). Geraadpleegd april 2017 via: <http://www.zorgvoorbeter.nl/ouderenzorg/Continentie-Praktijk-Signaleren.html>.
- Geen wetenschappelijke literatuur gevonden.

Arbocheck

- Arbocheck in de Thuiszorg. Een werkpakket om zelf mee aan de slag te gaan in het kader van de Praktijkregels (2004). Convenantpartijen Arbeidsomstandigheden Thuiszorg.

Barthel index

- Balu (2009). Differences in psychometric properties, cut-off scores, and outcomes between the Barthel Index and Modified Rankin Scale in pharmacotherapybased stroke trials: systematic literature review. *Curr Med Res Opin*;25(6):1329-41.
- Dawson (2008). A structured review of outcome measures used for the assessment of rehabilitation interventions for spinal cord injury. *Spinal Cord*;46:768–780
- Death e.a. (2009). Selection of outcome measures in lower extremity amputation rehabilitation: ICF activities. *Dis Rehab*;31(18):1455-1473,
- de Haan e.a. (1993). Klinimetrische evaluatie van de Barthel Index: een maat voor beperkingen in het dagelijks functioneren. *Ned Tijdschr Geneesk*;137(18):917-21.
- Hartigan (2007). A comparative review of the Katz ADL and the Barthel Index in assessing the activities of daily living of older people. *Int J Older People Nurs*;2(3):204-12.
- Post e.a. (1995). Nederlandse interviewversie van de Barthel-index onderzocht bij dwarslaesiepatiënten. *Ned Tijdschr Geneesk*;139(27):1376-1380.
- Sainsbury e.a (2005). Reliability of the Barthel Index when used with older people. *Age Ageing*;34(3):228-32.

BasisZorglijst - Continentiemodule

- BasisZorgLijst - 24 uurs registratie - Continentiemodule Verbetertraject (2008). LOCOmotion t.b.v. Verbetertraject Continentie VILANS / Zorg voor Beter. Geraadpleegd april 2017 via: <http://www.zorgvoorbeter.nl/ouderenzorg/Continentie-Leren-Incontinentiezorg-evaluieren.html>.
- Geen wetenschappelijke literatuur gevonden.

Beck depression inventory (BDI)

- Copyright, alleen tegen betaling beschikbaar. Vragenlijst te vinden in: Handleiding bouwstenen zorgpaden GGZ (2012). Coproductie van het Trimbos-instituut en het ROS-netwerk.
- Lako e.a. (2012). A systematic review of instruments to measure depressive symptoms in patients with schizophrenia. *J Affect Disord*;140(1):38-47.
- Stockings e.a. (2015). Symptom screening scales for detecting major depressive disorder in children and adolescents: a systematic review and meta-analysis of reliability, validity and diagnostic utility. *J Affect Disord*;15;174:447-63.
- Vodermaier e.a. (2009). Screening for emotional distress in cancer patients: a systematic review of assessment instruments. *J Natl Cancer Inst*. 2009 Nov 4;101(21):1464-88.
- Wang & Gorenstein (2013). Psychometric properties of the Beck Depression Inventory-II: a comprehensive review. *Rev Bras Psiquiatr*;35(4):416-31.

BEM (Beoordeling beheer Eigen Medicatie)

- Procedure Beoordeling beheer Eigen Medicatie (BEM) van en door thuiszorgorganisaties (2016). Instituut voor Verantwoord Medicijngebruik.
- Geen wetenschappelijke literatuur gevonden.

Beroepscode Verpleegkundigen en Verzorgenden

- Beroepscode van Verpleegkundigen en Verzorgenden (2015). GMV vakorganisatie voor christenen, CNV Zorg & Welzijn, FNV Zorg & Welzijn, HCF, NU'91, RMU Sector Gezondheidszorg en Welzijn 'Het Richtsnoer', V&VN.

Besluitvorming in de palliatieve fase

- Verbetertraject, integratie van richtlijnen Palliatieve Zorg. <https://www.iknl.nl/palliatieve-zorg/verbetertrajecten/besluitvorming-palliatieve-fase>.

Bradenschaal

- Braden & Maklebust (2005). Preventing pressure ulcers with the Braden scale: an update on this easy-to-use tool that assesses a patient's risk. *Am J Nurs*;105(6):70-2.
- Halfens e.a. (2000). Validity and reliability of the Braden scale and the influence of other risk factors: a multi-centre prospective study. *Int J Nurs Stud*;37(4):313-9.
- Kottner e.a. (2009). A systematic review of interrater reliability of pressure ulcer classification systems. *J Clin Nurs*;18(3):315-36.
- Kottner e.a. (2009). An interrater reliability study of the assessment of pressure ulcer risk using the Braden scale and the classification of pressure ulcers in a home care setting. *Int J Nurs Stud*;46(10):1307-12.
- Kring (2007). Reliability and validity of the Braden Scale for predicting pressure ulcer risk. *J Wound Ostomy Continence Nurs*;34(4):399-406.
- Pancorbo-Hidalgo e.a. (2006). Risk assessment scales for pressure ulcer prevention: a systematic review. *J Adv Nurs*;54(1):94-110.
- de Souza e.a. (2010). Predictive validity of the Braden Scale for Pressure Ulcer Risk in elderly residents of long-term care facilities. *Geriatr Nurs*;31(2):95-104.

Bristol Stoelgangschaal

- Blake e.a. (2016). Validity and reliability of the Bristol Stool Form Scale in healthy adults and patients with diarrhoea-predominant irritable bowel syndrome. *Aliment Pharmacol Ther*;44(7):693-703.
- Fallon e.a. (2008). A systematic review of psychometric evidence and expert opinion regarding the assessment of faecal incontinence in older community-dwelling adults. *Int J Evid Based Healthc*;6(2):225-59.
- Lewis & Heaton (1997). Stool form scale as a useful guide to intestinal transit time. *Scand J Gastroenterol*;32(9):920-4.

Care2App Omaha

- Zie Omaha system.

Caregiver Reaction Assessment (CRA)

- Harkness e.a. (2007). Measurement of the caregiving experience in caregivers of persons living with heart failure: a review of current instruments. *J Card Fail*;13(7):577-87.
- Nijboer e.a. (1999). Measuring both negative and positive reactions to giving care to cancer patients: psychometric qualities of the Caregiver Reaction Assessment (CRA). *Soc Sci Med*;48:1259-69.
- Post e.a. (2007). Reproducibility of the Caregiver Strain Index and the Caregiver Reaction Assessment in partners of stroke patients living in the Dutch community. *Clin Rehabil*;21(11):1050-5.
- Stephan e.a. (2013). Validity, reliability, and feasibility of the German version of the Caregiver Reaction Assessment scale (G-CRA): a validation study. *Int Psychogeriatr*;25(10):1621-8.
- Van Durme e.a. (2012). Tools for measuring the impact of informal caregiving of the elderly: a literature review. *Int J Nurs Stud*;49(4):490-504.
- Van Exel e.a. (2004). Instruments for assessing the burden of informal caregiving for stroke patients in clinical practice: a comparison of CSI, CRA, SCQ and self-rated burden. *Clin Rehabil*;18(2):203-14.

Caregiver Strain Index (CSI)

- Hudson e.a. (2010). A systematic review of instruments related to family caregivers of palliative care patients. *Palliative medicine*;24(7):656-668.
- Post e.a. (2007). Reproducibility of the Caregiver Strain Index and the Caregiver Reaction Assessment in partners of stroke patients living in the Dutch community. *Clin Rehabil*;21(11):1050-5.
- Robinson (1983). Validation of a Caregiver Strain Index. *J Gerontol*;38(3):344-348. *Volledige tekst van artikel niet online te vinden.*
- Van Durme e.a. (2012). Tools for measuring the impact of informal caregiving of the elderly: a literature review. *Int J Nurs Stud*;49(4):490-504.
- Van Exel e.a. (2004). Instruments for assessing the burden of informal caregiving for stroke patients in clinical practice: a comparison of CSI, CRA, SCQ and self-rated burden. *Clin Rehabil*;18(2):203-14.
- Whalen & Buchholz. (2009). The reliability, validity and feasibility of tools used to screen for caregiver burden: a systematic review. *JBI Libr Syst Rev*;7(32):1373-1430.

Carpenito

- Carpenito (2012). *Zakboek Verpleegkundige Diagnosen*. Noordhoff Uitgevers Groningen/Houten, 4^e druk, vertaling van de 13e editie van *Handbook of Nursing Diagnosis*.

CSDD Cornell Scale for Depression in Dementia

- Goodarzi e.a. (2017). Depression Case Finding in Individuals with Dementia: A Systematic Review and Meta-Analysis. *J Am Geriatr Soc*;65(5):937-948.
- Knapskog e.a. (2011). A comparison of the validity of the Cornell Scale and the MADRS in detecting depression among memory clinic patients. *Dement Geriatr Cogn Disord*;32(4):287-94.
- Kørner e.a. (2006). The Geriatric Depression Scale and the Cornell Scale for Depression in Dementia. A validity study. *Nord J Psychiatry*;60(5):360-4.
- Kørner e.a. (2007). Rating scales for depression in the elderly: external and internal validity. *J Clin Psychiatry*;68(3):384-9.
- Leontjevas e.a. (2009). The Montgomery-Asberg Depression Rating Scale and the Cornell Scale for Depression in Dementia: a validation study with patients exhibiting early-onset dementia. *Am J Geriatr Psychiatry*;17(1):56-64.
- Perrault e.a. (2000). Review of outcome measurement instruments in Alzheimer's disease drug trials: psychometric properties of behavior and mood scales. *J Geriatr Psychiatry Neurol*;13(4):181-96.

Communicatie bij dementie

- Communicatie bij dementie. Tien tips voor de communicatie met mensen met dementie. *Academische werkplaats Dementie*, 2010.

Competentielijst ondersteuning zelfmanagement

- van Hooft e.a. (2015). Ondersteunen van zelfmanagement: wat vraagt dit van verpleegkundigen? Ned Tijdschr Ev Based Pract;10(1):17-20.

Complexiteitsscorelijst, cliënttoewijzingsformulier en kritische signalenlijst

- Complexiteitsscorelijst, cliënttoewijzingsformulier en kritische signalenlijst (2005). Stichting Thuiszorg Maasmond/Hogeschool InHolland.
- Geen wetenschappelijke literatuur gevonden.

De mond niet vergeten

- <https://www.demondnievergeten.nl/screenings-en-verwijsinstrument-thuiszorg/>
- Geen wetenschappelijke literatuur gevonden.

DENWIS

- Douw e.a. (2017). Capturing early signs of deterioration: the dutch-early-nurse-worry-indicator-score and its value in the Rapid Response System. J Clin Nurs;26(17-18):2605-2613.
- Douw e.a. (2016). Nurses' 'worry' as predictor of deteriorating surgical ward patients: A prospective cohort study of the Dutch-Early-Nurse-Worry-Indicator-Score. Int J Nurs Stud;59:134-40.

Disability Rating Index

- Grotle (2012). Reliability and construct validity of self-report questionnaires for patients with pelvic girdle pain. Phys Ther;92(1):111-23.
- Parsons (2015). Measurement properties of the Disability Rating Index in patients undergoing hip replacement. Rheumatology (Oxford). 2015 Jan;54(1):64-71.
- Salén e.a. (1994). The Disability Rating Index: an instrument for the assessment of disability in clinical settings. J Clin Epidemiol;47(12):1423-3. Full text niet beschikbaar.
- Wittboldt e.a. (2016). Reliability of two questionnaires on physical function in patients with stable coronary artery disease. Eur J Cardiovasc Nurs;15(2):142-9.

DOLOPLUS-2

- Chen e.a. (2010). Validating nurses' and nursing assistants' report of assessing pain in older people with dementia. J Clin Nurs;19(1-2):42-52.
- Hølen e.a. (2007). Doloplus-2, a valid tool for behavioural pain assessment? BMC Geriatr;19;7:29.
- Neville e.a. (2014). A psychometric evaluation of three pain rating scales for people with moderate to severe dementia. Pain Manag Nurs. 2014 Dec;15(4):798-806.
- Smith e.a. (2013). Assessment of people with cognitive impairment and hip fracture: a systematic review and meta-analysis. Arch Gerontol Geriatr;57(2):117-26.
- Zwakhalen e.a. (2006). The psychometric quality and clinical usefulness of three pain assessment tools for elderly people with dementia. Pain;15;126(1-3):210-20.
- Zwakhalen e.a. (2006). Pain in elderly people with severe dementia: a systematic review of behavioural pain assessment tools. BMC Geriatr. 2006 Jan 27;6:3.

DOS Delier observatieschaal

- Grover (2012). Assessment scales for delirium: A review. World J Psychiatry;22;2(4):58-70.
- Timmers (2004). Een overzicht van beoordelingsschalen voor delier. Tijdschr Gerontol Geriatr;35(1):5-14.
- van Velthuijsen (2016). Psychometric properties and feasibility of instruments for the detection of delirium in older hospitalized patients: a systematic review. Int J Geriatr Psychiatry;31(9):974-89.

Duaal procesmodel

- Moorse (2015). Rouw mag er zijn. J Soc Interv: Theory Practice;24:45-57.
- Lund e.a. (2010). Experiences and Early Coping of Bereaved Spouses/Partners in an Intervention Based on the Dual Process Model (DPM). Omega; 61(4): 291–313.
- Richardson (2007). A dual process model of grief counseling: findings from the Changing Lives of Older Couples (CLOC) study (abstract only). J Gerontol Soc Work. 2007;48(3-4):311-29.
- Stroebe & Schut (1999). The dual process model of coping with bereavement - Rationale and description. Death Studies;23:197-224.
- Stroebe & Schut (2010). The Dual Process Model of Coping with Bereavement: A Decade on. OMEGA;61:273-289.

Easycare Tweetraps Ouderen Screening (TOS)

- Brandão e.a. (2017). Reliability and validity of the EASYCare-2010 Standard to assess elderly people in Portuguese Primary Health Care. Aten Primaria;S0212-6567(16)30271-2.
- Craig (2015). Systematic review of EASY-care needs assessment for community-dwelling older people. Age Ageing;44:559–565.
- Keiren e.a. (2014). Feasibility evaluation of a stepped procedure to identify community-dwelling frail older people in general practice. A mixed methods study. Eur J Gen Pract;20(2):107-13.
- Van Kempen e.a. (2013). Development of an instrument for the identification of frail older people as a target population for integrated care. Br J Gen Pract;63(608):e225-31.
- Van Kempen e.a. (2014). Construct validity and reliability of a two-step tool for the identification of frail older people in primary care. J Clin Epidemiol;67(2):176-83.
- Van Kempen e.a. (2015). Predictive validity of a two-step tool to map frailty in primary care. BMC Medicine; 13:287.

- Morley e.a. (2017). Integrated Care: Enhancing the Role of the Primary Health Care Professional in Preventing Functional Decline: A Systematic Review. *J Am Med Dir Assoc*;18(6):489-494.

Ecogram

- Hartman (1995). Diagrammatic assessment of family relationships. *Fam in Soc*;111-122.
- Ray & Street (2005). Ecomapping: an innovative research tool for nurses. *J Adv Nurs*;50(5):545-52.

EDIZ mantelzorgscan

- Pot, e.a. (1995). Ervaren druk door informele zorg; constructie van een schaal. *Tijdschr Gerontol Geriatr*;1995;26:214-219.
Volledige tekst van artikel niet online te vinden.

EDIZ Plus mantelzorgscan

- De Boer e.a. (2012). Self perceived burden from informal care: construction of the EDIZ-plus. *Tijdschr Gerontol Geriatr*;43(2):77-88.

Edmonton Functional Assessment Tool

- Gardiner & Samuel (2014). Evaluation of the Edmonton Functional Assessment Tool (EFAT2) within palliative care: a pilot study. *Aging Clin Exp Res*;26(3):293-8.
- Kaasa (1997). The Edmonton Functional Assessment Tool: preliminary development and evaluation for use in palliative care. *J Pain Symptom Manage*;13(1):10-9.
- Kaasa & Wessel (2001). The Edmonton Functional Assessment Tool: further development and validation for use in palliative care. *J Palliat Care*;17(1):5-11. *Volledige tekst van artikel niet online te vinden.*
- Kaasa e.a. (2000). Inter-rater reliability of formally trained and self-trained raters using the Edmonton Functional Assessment Tool. *Palliat Med*;14(6):509-17.

Edmonton Symptom Assessment System (ESAS) / Utrecht Symptoom Dagboek (USD)

- Aktas e.a. (2015). The psychometric properties of cancer multisymptom assessment instruments: a clinical review. *Support Care Cancer*;23(7):2189-202.
- Chang e.a. (2000). Validation of the Edmonton Symptom Assessment Scale. *Cancer*;88(9):2164-71. Full text niet beschikbaar.
- De Graaf (2017). Symptom intensity of hospice patients: A longitudinal analysis of concordance between patients' and nurses' outcomes. *J Pain Symptom Manage*; pii: S0885-3924(17)30469-4.
- Hannon e.a. (2015). Modified Edmonton Symptom Assessment System including constipation and sleep: validation in outpatients with cancer. *J Pain Symptom Manage*;49(5):945-52.
- Richardson & Jones (2009). A review of the reliability and validity of the Edmonton Symptom Assessment System. *Curr Oncol*; 16(1): 55.
- Watanabe e.a. (2012). The Edmonton Symptom Assessment System, a proposed tool for distress screening in cancer patients: development and refinement. *Psychooncology*;21(9):977-85. Full text niet beschikbaar.
- Zweers e.a. (2017). The predictive value of symptoms for anxiety in hospice inpatients with advanced cancer. *Palliat Support Care*:1-6.

Eenzaamheidsschaal

- De Jong Gierveld & van Tilburg (2008). De ingekorte schaal voor algemene, emotionele en sociale eenzaamheid. *Tijdschr Gerontol Geriatr*;39:1,4-15.
- De Jong Gierveld & van Tilburg (2010). The De Jong Gierveld short scales for emotional and social loneliness: tested on data from 7 countries in the UN generations and gender surveys. *Eur J Ageing*;7(2):121-130.
- Uysal-Bozkir e.a. (2017). Translation and Validation of the De Jong Gierveld Loneliness Scale Among Older Migrants Living in the Netherlands. *J Gerontol B Psychol Sci Soc Sci*;72(1):109-119.

EFIP Evaluative Frailty Index for Physical activity

- de Vries e.a. (2013). Evaluative frailty index for physical activity (EFIP): a reliable and valid instrument to measure changes in level of frailty. *Phys Ther*;93(4):551-61.

eHealth indiceren in de wijk

- eHealth indiceren in de wijk. Windesheim, 2017 (in ontwikkeling).

eOverdracht

- Duijvendijk (2014). eOverdracht in de praktijk: ervaringen en tips van gebruikers. Nictiz.

EQ-5D-5L

- EuroQol (2015). EQ-5D-5L User Guide. Basic information on how to use the EQ-5D-5L instrument.

EWS (Early Warning Score)

- Alam e.a. (2014). The impact of the use of the Early Warning Score (EWS) on patient outcomes: a systematic review. *Resuscitation*;85(5):587-94.
- Liljequist & Christensen (2016). Early warning score predicts acute mortality in stroke patients. *Acta Neurol Scand*;133(4):261-7.
- Stark e.a. (2015). An Early Warning Score Predicts Risk of Death after In-hospital Cardiopulmonary Arrest in Surgical Patients. *Am Surg*;81(10):916-21.
- Williams e.a. (2016). The ability of early warning scores (EWS) to detect critical illness in the prehospital setting: A systematic review. *Resuscitation*;102:35-43.

Factsheet Zorg voor kinderen met een intensieve zorgvraag

- <https://www.rijksoverheid.nl/documenten/brochures/2016/11/03/factsheet-zorg-voor-kinderen-met-een-intensieve-zorgvraag-algemeen>, geraadpleegd augustus 2017.

Fear-Avoidance Beliefs Questionnaire

- Bishop (2007). Health care practitioners' attitudes and beliefs about low back pain: a systematic search and critical review of available measurement tools. *Pain*;132(1-2):91-101.
- Chapman (2011). Evaluating common outcomes for measuring treatment success for chronic low back pain. *Spine (Phila Pa 1976)*;36(21 Suppl):S54-68.
- Grotle (2012). Reliability and construct validity of self-report questionnaires for patients with pelvic girdle pain. *Phys Ther*;92(1):111-23.
- Lundberg (2011). Pain-related fear: a critical review of the related measures. *Pain Res Treat*:494196.
- Mateen (2017). Systematic Review of Health-Related Work Outcome Measures and Quality Criteria-Based Evaluations of Their Psychometric Properties. *Arch Phys Med Rehabil*;98(3):534-560.

Fecal Incontinence Quality of Life Scale

- Bols e.a. (2013). Responsiveness and interpretability of incontinence severity scores and FIQL in patients with fecal incontinence: a secondary analysis from a randomized controlled trial. *Int Urogynecol J*;24(3):469-78.
- Fallon e.a. (2008). A systematic review of psychometric evidence and expert opinion regarding the assessment of faecal incontinence in older community-dwelling adults. *Int J Evid Based Healthc*;6(2):225-59.
- 't Hoen e.a. (2017). The fecal incontinence quality of life scale (FIQL) and fecal incontinence severity index (FISI): Validation of the Dutch versions. *Neurourol Urodyn*;36(3):710-715.
- Rockwood e.a. (2000). Fecal Incontinence Quality of Life Scale: quality of life instrument for patients with fecal incontinence. *Dis Colon Rectum*;43(1):9-16; discussion 16-7.
- Trajanovska & Catto-Smith (2005). Quality of life measures for fecal incontinence and their use in children. *J Gastroenterol Hepatol*;20(6):919-28.

Formulier mantelzorg

- Formulier Mantelzorg in het zorg(leef)plan, Toelichting en instructie (2012). Co-productie van de Brabantse Raad voor Informele Zorg, Expertisecentrum Mantelzorg, Integraal Kankercentrum Zuid en Vilans.

GDS Geriatrische depressie schaal

- Allen & Annells (2009). A literature review of the application of the Geriatric Depression Scale, Depression Anxiety Stress Scales and Post-traumatic Stress Disorder Checklist to community nursing cohorts. *J Clin Nurs*;18(7):949-59.
- Burton & Tyson (2015). Screening for mood disorders after stroke: a systematic review of psychometric properties and clinical utility. *Psychol Med*;45(1):29-49.
- De Bruyne e.a. (2009). Is the geriatric depression scale a reliable screening tool for depressive symptoms in elderly patients with cognitive impairment? *Int J Geriatr Psychiatry*;24(6):556-62.
- Goodarzi e.a. (2017). Depression Case Finding in Individuals with Dementia: A Systematic Review and Meta-Analysis. *J Am Geriatr Soc*;65(5):937-948.
- Mitchell e.a. (2010). Diagnostic validity and added value of the Geriatric Depression Scale for depression in primary care. *J Affect Disord*;125(1-3):10-7.
- Kørner e.a. (2006). The Geriatric Depression Scale and the Cornell Scale for Depression in Dementia. A validity study. *Nord J Psychiatry*;60(5):360-4.
- Pocklington e.a. (2016). The diagnostic accuracy of brief versions of the Geriatric Depression Scale: a systematic review and meta-analysis. *Int J Geriatr Psychiatry*;31(8):837-57.
- Stiles & McGarrahian (1998). The Geriatric Depression Scale: A Comprehensive Review. *J of Clin Geropsychol*;4:89-111.
- Wancata (2006). The criterion validity of the Geriatric Depression Scale: a systematic review. *Acta Psychiatr Scand*;114(6):398-410.

Gebruikelijke Zorg

- Gebruikelijke zorg (-). Rijksoverheid. Geraadpleegd april 2017 via:
https://www.regelhulp.nl/bladeren/_/artikel/gebruikelijke-zorg/

General self-efficacy scale

- De Las Cuevas & Peñate (2015). Validation of the General Self-Efficacy Scale in psychiatric outpatient care. *Psicothema*;27(4):410-5.
- Kupst e.a. (2015). Assessment of stress and self-efficacy for the NIH Toolbox for Neurological and Behavioral Function. *Anxiety Stress Coping*;28(5):531-44.
- Machado e.a. (2016). Psychometric properties of Multidimensional Health Locus of Control - A and General Self-Efficacy Scale in civil servants: ELSA-Brasil Musculoskeletal Study (ELSA-Brasil MSK). *Braz J Phys Ther*;20(5):451-460.
- Nilsson e.a. (2015). Psychometric properties of the General Self-Efficacy Scale in Parkinson's disease. *Acta Neurol Scand*;132(2):89-96.
- Ohno e.a. (2017). Smallest detectable change and test-retest reliability of a self-reported outcome measure: Results of the Center for Epidemiological Studies Depression Scale, General Self-Efficacy Scale, and 12-item General Health Questionnaire. *J Eval Clin Pract*. [Epub ahead of print]
- Scholz e.a. (2002). Is General Self-Efficacy a universal construct? Psychometric Findings from 25 Countries. *Eur J Psychol Assessment*; 18(30): 242-251.

Genogram

- Een genogram maken: de generaties in beeld. Expertisecentrum Mantelzorg. Geraadpleegd april 2017 via:
<http://www.expertisecentrummantelzorg.nl/em/instrument-een-genogram-maken-de-generaties-in-beeld.html>
- Puskar (1996). Genogram: a useful tool for nurse practitioners. *J Psychiatr Ment Health Nurs*;3(1):55-60.
- Rogers e.a. (1990). Completion rate and reliability of the self-administered genogram (SAGE). *Fam Pract*;7(2):149-51.
Volledige tekst van artikel niet online te vinden.

- Rogers e.a. (1991). The SAGE-PAGE trial: do family genograms make a difference? *J Am Board Fam Pract*;4(5):319-26.

Gesprekskaart 'Baas over je eigen gezondheid'

- Wolters & Engels (2016). Gesprekskaart 'Baas over je eigen gezondheid'. Vilans en Kennisplein Chronische Zorg.

Gestructureerde Medicatie Anamnese

- Drenth-van Maanen e.a. (2011). Structured history taking of medication use reveals iatrogenic harm due to discrepancies in medication histories in hospital and pharmacy records. *J Am Geriatr Soc*;59(10):1976-7.
- Gestructureerde Medicatie Anamnese (2010). Afdeling Geriatrie en Expertisecentrum Pharmacotherapie bij Ouderen (PHOR) en afdeling Farmacie UMC Utrecht.
- Spee e.a. (2010). Gestructureerde medicatieanamnese bij opname, helpt een juist overzicht van de thuismedicatie te verkrijgen. *Ned tijdschr geneesk*;154:A904.

Gordon, 11 gezondheidspatronen

- Gordon (1982). Nursing Diagnosis: Process and Application. McGraw-Hill Inc.US, 3rd Edition.

Goed in gesprek met mantelzorgers

- <http://www.expertisecentrummantelzorg.nl/em.net?id=10037>
- Geen wetenschappelijke literatuur gevonden.

Groningen Activity Restriction Scale (GARS)

- Douglas e.a. (1995). The assessment of functional status in rheumatoid arthritis: a cross cultural, longitudinal comparison of the Health Assessment Questionnaire and the Groningen Activity Restriction Scale. *J Rheumatol*;22(10):1834-43. *Volledige tekst van artikel niet online te vinden.*
- Jansen e.a. (2010). Psychometric properties of questionnaires evaluating health-related quality of life and functional status in polytrauma patients with lower extremity injury. *J Trauma Manag Outcomes*;28:4:7.
- Kempen e.a. (1996). The assessment of disability with the Groningen Activity Restriction Scale. Conceptual framework and psychometric properties. *Soc Sci Med*;43(11):1601-10.
- Kempen e.a. (2012). Groningen Activiteiten Restrictie Schaal (GARS). Een handleiding. Research Institute SHARE, UMCG / Rijksuniversiteit Groningen, 2^e druk.
- Oude Voshaar (2011). Measurement properties of physical function scales validated for use in patients with rheumatoid arthritis: a systematic review of the literature. *Health Qual Life Outcomes*;7:9:99.
- Suurmeijer e.a. (1994). The Groningen Activity Restriction Scale for measuring disability: its utility in international comparisons. *Am J Public Health*;84(8):1270-3.
- Swinkels e.a. (2005). Reliability, validity and responsiveness of instruments to assess disabilities in personal care in patients with rheumatic disorders. A systematic review. *Clin Exp Rheumatol*;23(1):71-9.

Groningen Frailty Indicator (GFI)

- Apóstolo e.a. (2017). Predicting risk and outcomes for frail older adults: an umbrella review of frailty screening tools. *JBI Database System Rev Implement Rep*;15(4):1154-1208.
- Daniels e.a. (2012). The predictive validity of three self-report screening instruments for identifying frail older people in the community. *BMC Public Health*;23:12:69.
- Dent e.a. (2016). Frailty measurement in research and clinical practice: A review. *Eur J Intern Med*;31:3-10.
- Hoogendijk e.a. (2013). The identification of frail older adults in primary care: comparing the accuracy of five simple instruments. *Age Ageing*;42(2):262-5.
- Metzelthin e.a. (2010). The psychometric properties of three self-report screening instruments for identifying frail older people in the community. *BMC Public Health*;31:10:176.
- Peters e.a. (2012). Measurement properties of the Groningen Frailty Indicator in home-dwelling and institutionalized elderly people. *J Am Med Dir Assoc*;13(6):546-51.
- Peters e.a. (2015). Construct validity of the Groningen Frailty Indicator established in a large sample of home-dwelling elderly persons: Evidence of stability across age and gender. *Exp Gerontol*;69:129-41.
- Steverink e.a. (2001). Measuring frailty: Developing and testing the GFI (Groningen frailty indicator). [Geen fulltext of abstract kunnen vinden]
- Sutorius e.a. (2016). Comparison of 10 single and stepped methods to identify frail older persons in primary care: diagnostic and prognostic accuracy. *BMC Fam Pract*;3:17:102.

HADS Hospital Anxiety and Depression scale

- Bjelland e.a. (2002). The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *J Psychosom Res*;52(2):69-77.
- Burton & Tyson (2015). Screening for mood disorders after stroke: a systematic review of psychometric properties and clinical utility. *Psychol Med*;45(1):29-49.
- Litster e.a. (2016). Screening Tools for Anxiety in People with Multiple Sclerosis: A Systematic Review. *Int J MS Care*;18(6):273-281.
- Petterson e.a. (2015). Which instruments to support diagnosis of depression have sufficient accuracy? A systematic review. *Nord J Psychiatry*;69(7):497-508.
- Vodermaier (2011). Accuracy of the Hospital Anxiety and Depression Scale as a screening tool in cancer patients: a systematic review and meta-analysis. *Support Care Cancer*;19(12):1899-908.

Handleiding Zorgen voor Zelfzorg

- Handleiding Zorgen voor Zelfzorg: Tips en ervaringen uit de werkplaats zelfmanagement (2015). Vilans, Zelfzorg Ondersteund (ZO!), InEen in samenwerking met Eerstelijns Zorggroep Haaglanden (ELZHA), SGE uit Eindhoven en Zorggroep Almere.

Handreiking Kwetsbare Ouderen

- Pel-Littel e.a. (2014). Kwetsbare Ouderen. Handreiking voor professionals & multidisciplinaire teams in de eerste lijn. Ontwikkeld door Vilans e.a.

Handreiking zelfredzaamheid voor wijkverpleegkundigen

- Jansen (2013). Handreiking zelfredzaamheid voor wijkverpleegkundigen. Vilans.

HDRS depressie schaal

- Bagby e.a. (2004). The Hamilton Depression Rating Scale: has the gold standard become a lead weight? Am J Psychiatry;161(12):2163-77.
- Berg e.a. (2009). Assessment of depression after stroke: a comparison of different screening instruments. Stroke;40(2):523-9.
- Goodarzi e.a. (2017). Depression Case Finding in Individuals with Dementia: A Systematic Review and Meta-Analysis. J Am Geriatr Soc;65(5):937-948.
- Meader e.a. (2014). Screening for poststroke major depression: a meta-analysis of diagnostic validity studies. J Neurol Neurosurg Psychiatry;85(2):198-206.
- Trajković e.a. (2011). Reliability of the Hamilton Rating Scale for Depression: a meta-analysis over a period of 49 years. Psychiatry Res;30:189(1):1-9.
- Belgian Screening Tools, HDRS bespreking betrouwbaarheid. Geraadpleegd mei 2017 via: http://www.best.ugent.be/BEST3_NL/nederlands_schalen_bespreking_schalen_hdrs_hdrs4.html

Huidproblemen bij incontinentie

- E-learning - Huidproblemen bij incontinentie. Geraadpleegd april 2017 via: <http://www.vilans.nl/nieuwsoverzicht-elearning-huidproblemen-incontinentie.html>.

ICF checklist

- Dernek e.a. (2015). The utility of the International Classification of Functioning, Disability and Health checklist for evaluating disability in a community-dwelling geriatric population sample. Int J Rehabil Res;38(2):144-55.
- Müller-Staub e.a. (2007). Meeting the criteria of a nursing diagnosis classification: Evaluation of ICNP, ICF, NANDA and ZEFP. Int J Nurs Stud;44(5):702-13.
- Stallinga e.a. (2014). Functioning assessment vs. conventional medical assessment: a comparative study on health professionals' clinical decision-making and the fit with patient's own perspective of health. J Clin Nurs;23(7-8):1044-54.

Indicatiwegwijzer

- Geen wetenschappelijke literatuur gevonden.

Indiceren en zorgtoewijzing

- Cursus Indiceren en Zorgtoewijzing voor Wijkverpleegkundigen. Zuyd Professional. <https://www.zuydprofessional.nl/opleidingen/indiceren-en-zorgtoewijzing-voor-wijkverpleegkundigen>

Informatiekaart informele en formele zorg

- Informatiekaart informele en formele zorg. Aandacht voor iedereen. Geraadpleegd april 2017 via: www.aandachtvoriedereen.nl.

INTERMED

- Huyse e.a. (1999). "INTERMED": a method to assess health service needs. I. Development and reliability. Gen Hosp Psychiatry;21(1):39-48.
- De Jonge e.a. (2005). Operationalization of biopsychosocial case complexity in general health care: the INTERMED project. Aust N Z J Psychiatry;39(9):795-9.
- Latour e.a. (2007). A method to provide integrated care for complex medically ill patients: the INTERMED. Nurs Health Sci;9(2):150-7.
- Peters e.a. (2013). Development and measurement properties of the self assessment version of the INTERMED for the elderly to assess case complexity. J Psychosom Res;74(6):518-22.
- Van Reedt Dordtland e.a. (2017). Assessment of Biopsychosocial Complexity and Health Care Needs: Measurement Properties of the INTERMED Self-Assessment Version. Psychosom Med;79(4):485-492.
- Stiefel e.a. (1999). "INTERMED": a method to assess health service needs. II. Results on its validity and clinical use. Gen Hosp Psychiatry;21(1):49-56.
- Stiefel e.a. (2006). Operationalizing integrated care on a clinical level: the INTERMED project. Med Clin North Am;90(4):713-58.

Inventarisatie zelfzorg mondzorg

- Inventarisatie zelfzorg mondzorg (2007). Uit: Richtlijn Mondzorg voor zorgafhankelijke cliënten in verpleeghuizen, NVVA (nu Verenso), bijlage 2a.

IPQ-K Illness Perception Questionnaire-Kort (IPQ-K)

- Broadbent e.a. (2015). A systematic review and meta-analysis of the Brief Illness Perception Questionnaire. Psychol Health. 2015;30(11):1361-85.
- Leysen e.a. (2015). Clinimetric properties of illness perception questionnaire revised (IPQ-R) and brief illness perception questionnaire (Brief IPQ) in patients with musculoskeletal disorders: A systematic review. Man Ther;20(1):10-7.
- de Raaij e.a. (2012). Cross-cultural adaptation and measurement properties of the Brief Illness Perception Questionnaire-Dutch Language Version. Man Ther;17(4):330-5.

Is alles besproken?

- Vragenlijst 'Is alles besproken': behoud van kwaliteit van leven. Geraadpleegd april 2017 via: <http://www.beteroud.nl/ouderen/zorg-is-alles-besproken-vragenlijst.html> en <http://www.nuzo-utrecht.nl/Projecten-Best-Practices/NUZO-projecten-lopend>.

Katz ADL schaal

- Buurman e.a. (2011). Variability in measuring (instrumental) activities of daily living functioning and functional decline in hospitalized older medical patients: a systematic review. *J Clin Epidemiol*;64(6):619-27.
- Cabañero-Martínez e.a. (2009). The Spanish versions of the Barthel index (BI) and the Katz index (KI) of activities of daily living (ADL): a structured review. *Arch Gerontol Geriatr*;49(1):e77-84.
- Hartigan (2007). A comparative review of the Katz ADL and the Barthel Index in assessing the activities of daily living of older people. *Int J Older People Nurs*;2(3):204-12.
- Katz e.a. (1963). Studies of illness in the aged. The index of adl: a standardized measure of biological and psychosocial function. *Jama*;185:914-9.
- Roedl e.a. (2016). A systematic review and comparison of functional assessments of community-dwelling elderly patients. *J Am Assoc Nurse Pract*;28(3):160-9.

Kickprotocollen

- Alleen toegankelijk voor abonnees.

Kloktekening

- Bossers e.a. (2012). Recommended measures for the assessment of cognitive and physical performance in older patients with dementia: a systematic review. *Dement Geriatr Cogn Dis Extra*;2(1):589-609.
- Ehreke e.a. (2010). Is the Clock Drawing Test a screening tool for the diagnosis of mild cognitive impairment? A systematic review. *Int Psychogeriatr*;22(1):56-63.
- Elzen e.a. (2004). De diagnostische waarde van de kloktekening in een klinische setting. *Tijdschr gerontol geriatr*;35;90-104.
- Müller e.a. (2017). Increased Diagnostic Accuracy of Digital vs. Conventional Clock Drawing Test for Discrimination of Patients in the Early Course of Alzheimer's Disease from Cognitively Healthy Individuals. *Front Aging Neurosci*;9:101.
- Nair e.a. (2010). Clock drawing test ratings by dementia specialists: interrater reliability and diagnostic accuracy. *J Neuropsychiatry Clin Neurosci*;22(1):85-92.
- Pinto e.a. (2009). Literature review of the Clock Drawing Test as a tool for cognitive screening. *Dement Geriatr Cogn Disord*;27(3):201-13.

Kwaliteit van leven vragenlijst voor mensen met ademhalingsproblemen

- Maillé & Kaptein (1993). Kwaliteit van leven vragenlijst voor mensen met ademhalingsproblemen. Leiden Universitair Medisch Centrum.

Landelijke richtlijn spirituele zorg

- Spirituele zorg (2010). Landelijke richtlijn, Versie: 1.0. Agora werkgroep. <http://www.oncoline.nl/spirituele-zorg>

Lastige gesprekken voeren

- Geraadpleegd mei 2017 via: <http://www.zorgvoorbeter.nl/ouderenzorg/Communiceren-Materialen-voor-medewerkers.html>
- Geen wetenschappelijke literatuur gevonden.

Lastmeter

- Donovan e.a. (2014). Validation of the distress thermometer worldwide: state of the science. *Psychooncology*;23(3):241-50.
- Tuinman e.a. (2008). Screening and referral for psychosocial distress in oncologic practice: use of the Distress Thermometer. *Cancer*;15;113(4):870-8.
- van Oers e.a. (2017). Dutch normative data and psychometric properties for the Distress Thermometer for Parents. *Qual Life Res*;26(1):177-182.
- Vodermaier e.a. (2009). Screening for emotional distress in cancer patients: a systematic review of assessment instruments. *J Natl Cancer Inst*. 2009 Nov 4;101(21):1464-88.
- Van overige bekijken artikelen/reviews (o.a. Miller (2013), Mitchel (2010), Stewart-Knight (2012), Snowden (2011)) geen full-text beschikbaar.

Lawton & Brody IADL vragenlijst

- Buurman e.a. (2011). Variability in measuring (instrumental) activities of daily living functioning and functional decline in hospitalized older medical patients: a systematic review. *J Clin Epidemiol*;64(6):619-27.
- Lawton & Brody (1969). Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist*;9(3):179-86.
- Roedl e.a. (2016). A systematic review and comparison of functional assessments of community-dwelling elderly patients. *J Am Assoc Nurse Pract*;28(3):160-9.
- Vittengl e.a. (2006). Comparative validity of seven scoring systems for the instrumental activities of daily living scale in rural elders. *Aging Ment Health*;10(1):40-7.

MADRS depressie schaal (Montgomery–Åsberg Depression Rating Scale)

- Fantino & Moore (2009). The self-reported Montgomery-Asberg Depression Rating Scale is a useful evaluative tool in Major Depressive Disorder. *BMC Psychiatry*;27;9:26.
- Furukawa (2010). Assessment of mood: guides for clinicians. *J Psychosom Res*;68(6):581-9.
- Goodarzi e.a. (2017). Depression Case Finding in Individuals with Dementia: A Systematic Review and Meta-Analysis. *J Am Geriatr Soc*;65(5):937-948.
- Knapskog e.a. (2011). A comparison of the validity of the Cornell Scale and the MADRS in detecting depression among memory clinic patients. *Dement Geriatr Cogn Disord*;32(4):287-94.

- Lako e.a. (2012). A systematic review of instruments to measure depressive symptoms in patients with schizophrenia. *J Affect Disord*;140(1):38-47.
- Leontjevas e.a. (2009). The Montgomery-Asberg Depression Rating Scale and the Cornell Scale for Depression in Dementia: a validation study with patients exhibiting early-onset dementia. *Am J Geriatr Psychiatry*;17(1):56-64.

MantelScan

- Kruijswijk (2012). Handreiking MantelScan. Samenstelling, organisatie en risicofactoren van een zorgnetwerk in kaart. Expertisecentrum Mantelzorg.
- Gebaseerd op bestaande instrumenten. Geen wetenschappelijke literatuur gevonden over de MantelScan als geheel.

Medicatieveiligheid achter de voordeur

- Medicatieveiligheid achter de voordeur. Werkmateriaal voor patiëntveiligheid in de eerste lijn (2010). Module is ontwikkeld door het Instituut voor Verantwoord Medicijngebruik voor en in samenwerking met het landelijke NEO/LVG patiëntveiligheidsprogramma 'Zorg voor Veilig'.

Mictiedagboek

- Bright e.a. (2014). Developing and validating the International Consultation on Incontinence Questionnaire bladder diary. *Eur Urol*;66(2):294-300.
- Bright e.a. (2011). Urinary diaries: evidence for the development and validation of diary content, format, and duration. *Neurourol Urodyn*;30(3):348-52.
- Brown e.a. (2003). Measurement characteristics of a voiding diary for use by men and women with overactive bladder. *Urology*;61(4):802-9.
- Martin e.a. (2006). Systematic review and evaluation of methods of assessing urinary incontinence. *Health Technol Assess*;10(6):1-132, iii-iv.

Mijnkwaliteitvanleven

- Hendrikx e.a. (2019). Measuring Population Health from a Broader Perspective: Assessing the My Quality of Life Questionnaire. *International J Integr Care*, 2019; 19(2): 7, 1–7.
- Beleidsrapportage 6 - Validatie van de vragenlijst van Mijnkwaliteitvanleven.nl. Mijnkwaliteitvanleven.nl, december 2018.

MMSE

- Bossers e.a. (2012). Recommended measures for the assessment of cognitive and physical performance in older patients with dementia: a systematic review. *Dement Geriatr Cogn Dis Extra*;2(1):589-609.
- Folstein e.a. (1975). "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*;12(3):189-198.
- Kok & Verhey (2002). Dutch translation of the Mini Mental State Examination (Folstein et al., 1975).
- Lopez e.a. (2005). Psychometric Properties of the Folstein Mini-Mental State Examination. *Assessment*;12:137.
- Paddick e.a. (2017). Cognitive screening tools for identification of dementia in illiterate and low-educated older adults, a systematic review and meta-analysis. *Int Psychogeriatr*;9:1-33.
- Pangman e.a. (2000). An Examination of Psychometric Properties of the Mini-Mental State Examination and the Standardized Mini-Mental State Examination: Implications for Clinical Practice. *Appl Nurs Res*;13:209-213

MNA-SF

- Bááth e.a. (2008). Interrater reliability using Modified Norton Scale, Pressure Ulcer Card, Short Form-Mini Nutritional Assessment by registered and enrolled nurses in clinical practice. *J Clin Nurs*;17(5):618-26.
- Guigoz (2006). The Mini Nutritional Assessment (MNA) review of the literature--What does it tell us? *J Nutr Health Aging*;10(6):466-85.
- Huhmann e.a. (2013). A self-completed nutrition screening tool for community-dwelling older adults with high reliability: a comparison study. *J Nutr Health Aging*;17(4):339-44.
- Phillips e.a. (2010). Nutritional screening in community-dwelling older adults: a systematic literature review. *Asia Pac J Clin Nutr*;19(3):440-9.
- Rubenstein (2001). Screening for undernutrition in geriatric practice: developing the short-form mini-nutritional assessment (MNA-SF). *J Gerontol A Biol Sci Med Sci*;56(6):M366-72.
- Skipper e.a. (2012). Nutrition screening tools: an analysis of the evidence. *JPEN J Parenter Enteral Nutr*;36(3):292-8.
- Vellas e.a. (2006). Overview of the MNA--Its history and challenges. *J Nutr Health Aging*;10(6):456-63.

Mobiliteits Protocol Toiletgang

- Mobiliteits Protocol Toiletgang. Zorg voor Beter (publicatiedatum onbekend).
- Geen wetenschappelijke literatuur gevonden.

MOCA Montreal Cognitive Assessment

- Burton & Tyson (2015). Screening for cognitive impairment after stroke: A systematic review of psychometric properties and clinical utility. *J Rehabil Med*;47(3):193-203.
- Ciesielska (2016). Is the Montreal Cognitive Assessment (MoCA) test better suited than the Mini-Mental State Examination (MMSE) in mild cognitive impairment (MCI) detection among people aged over 60? Meta-analysis. *Psychiatr Pol*;31;50(5):1039-1052.
- Isenberg-Grzeda e.a. (2017). A review of cognitive screening tools in cancer. *Curr Opin Support Palliat Care*;11(1):24-31.
- Ozer e.a. (2016). A systematic review of the diagnostic test accuracy of brief cognitive tests to detect amnestic mild cognitive impairment. *Int J Geriatr Psychiatry*;31(11):1139-1150.
- Tsoi e.a. (2015). Cognitive Tests to Detect Dementia: A Systematic Review and Meta-analysis. *JAMA Intern Med*;175(9):1450-8.
- Velayudhan e.a. (2014). Review of brief cognitive tests for patients with suspected dementia. *Int Psychogeriatr*;26(8):1247-62.

Mondstatuscorelijst

- Mondstatuscorelijst (2008). Afdeling Medische Oncologie, UMC Utrecht.
- Geen wetenschappelijke literatuur gevonden.

Mondzorg & de rol van verzorgenden en verpleegkundigen

- Mondzorg & de rol van verzorgenden en verpleegkundigen. Een samenvatting van de ‘Richtlijn Mondzorg voor zorgafhankelijke cliënten in verpleeghuizen’. NVVA, 2007.

Motiverende gesprekstechnieken zelfredzaamheid

- Motiverende gesprekstechnieken zelfredzaamheid (2014). Vilans.
- Geen wetenschappelijke literatuur gevonden.

MUST

- Anthony (2008). Nutrition screening tools for hospitalized patients. Nutr Clin Pract;23(4):373-82.
- Kruizenga (2010). The SNAQ(RC), an easy traffic light system as a first step in the recognition of undernutrition in residential care. J Nutr Health Aging;14(2):83-9.
- Phillips e.a. (2010). Nutritional screening in community-dwelling older adults: a systematic literature review. Asia Pac J Clin Nutr;19(3):440-9.
- Van Venrooij e.a. (2007). Quick-and-easy nutritional screening tools to detect disease-related undernutrition in hospital in- and outpatient setting: A systematic review of sensitivity and specificity. Eur J Clin Nutr Metab;2:21-37.
- Yaxley e.a. (2015). Identifying Malnutrition in an Elderly Ambulatory Rehabilitation Population: Agreement between Mini Nutritional Assessment and Validated Screening Tools. Healthcare (Basel);11;3(3):822-9.

NANDA NIC NOC

- Azzolin e.a. (2013). Effectiveness of nursing interventions in heart failure patients in home care using NANDA-I, NIC, and NOC. Appl Nurs Res;26(4):239-44.
- Boldt e.a. (2010). Combined application of the International Classification of Functioning, Disability and Health and the NANDA-International Taxonomy II. J Adv Nurs;66(8):1885-98.
- Bulechek e.a. (2007). Nursing Interventions Classification (Nic). Elsevier - Health Sciences Division, 5th revised edition.
- Eda e.a. (2014). Applicability of the NANDA-I and Nursing Interventions Classification taxonomies to mental health nursing practice. Int J Nurs Knowl;25(3):168-72.
- Moorhead e.a. (2012). Nursing Outcomes Classification (NOC), Measurement of Health Outcomes. Elsevier, 5th Edition.
- Müller-Staub e.a. (2007). Improved quality of nursing documentation: results of a nursing diagnoses, interventions, and outcomes implementation study. Int J Nurs Terminol Classif;18(1):5-17.
- Müller-Staub e.a. (2007). Meeting the criteria of a nursing diagnosis classification: Evaluation of ICNP, ICF, NANDA and ZEFP. Int J Nurs Stud;44(5):702-13.
- Rabelo-Silva et al (2017). Advanced Nursing Process quality: Comparing the International Classification for Nursing Practice (ICNP) with the NANDA-International (NANDA-I) and Nursing Interventions Classification (NIC). J Clin Nurs;26(3-4):379-387.
- Varsi & Ruland (2009). Congruence between nursing problems in nursing care plans and NANDA nursing diagnoses. Stud Health Technol Inform;146:778-9.

Nationale Kernset Patiëntproblemen

- Kieft (2015). Nationale Kernset - Beroepsstandaard voor het vastleggen van zorggegevens. V&VN. Nationale Kernset Patiëntproblemen, Geraadpleegd mei 2017 via: <https://www.nictiz.nl/terminologiecentrum/referentielijsten/nationale-kernset/nationale-kernset-patiëntproblemen>.
- Gebaseerd op kennis van verpleegkundigen en verzorgenden (totaal 985) uit alle zorgsectoren: ziekenhuis-, thuiszorg-, verpleeghuis- en verzorgingssector, geestelijke gezondheidszorg, verstandelijk gehandicaptenzorg en huisartsenpraktijk.

Niet Pluis Index

- Niet pluis index (2007). Welnis Preventie.
- Geen wetenschappelijke literatuur gevonden.

Normenkader indiceren en organiseren van zorg

- V&VN Normen voor indiceren en organiseren van verpleging en verzorging in de eigen omgeving (2014).

Nortonschaal

- Anthony e.a. (2008). Norton, Waterlow and Braden scores: a review of the literature and a comparison between the scores and clinical judgement. J Clin Nurs;17(5):646-53.
- Anthony e.a. (2010). Do risk assessment scales for pressure ulcers work? J Tissue Viability;19(4):132-6.
- Balzer e.a. (2007). The Norton, Waterlow, Braden, and Care Dependency Scales: comparing their validity when identifying patients' pressure sore risk. J Wound Ostomy Continence Nurs;34(4):389-98.
- Pancorbo-Hidalgo e.a. (2006). Risk assessment scales for pressure ulcer prevention: a systematic review. J Adv Nurs;54(1):94-110.
- Park e.a. (2016). Predictive Validity of Pressure Ulcer Risk Assessment Tools for Elderly: A Meta-Analysis. West J Nurs Res;38(4):459-83.
- Park & Lee (2016). Assessing Predictive Validity of Pressure Ulcer Risk Scales- A Systematic Review and Meta-Analysis. Iran J Public Health;45(2):122-33.
- Wang e.a. (2015). Inter-rater reliability of three most commonly used pressure ulcer risk assessment scales in clinical practice. Int Wound J;12(5):590-4.

Observatielijst dementie (OLD)

- Hopman-Rock e.a. (2001). Development and validation of the Observation List for early signs of Dementia (OLD). Int J Geriatr Psychiatry. 2001 Apr;16(4):406-14.

- Hopman-Rock e.a. (2001). Signalering van beginnende Alzheimer-dementie in de huisartspraktijk: Ontwikkeling en validering van de observatielijst voor vroege symptomen van dementie (OLD). *Tijdschr Gerontol Geriatr*;2;32:74-81.

Observatielijst voor psychosociale problematiek bij ouderen (OLP)

- Tak e.a. (2016). Development and preliminary validation of an Observation List for detecting mental disorders and social Problems in the elderly in primary and home care (OLP). *Int J Geriatr Psychiatry*;31(7):755-64.

Omaha System

- Radhakrishnan e.a. (2016). Effective Teaching-Learning Strategies for the Omaha System. *Home Healthc Now*;34(2):86-92.
- Ker e.a. (2016). Feasibility of Using the Omaha System for Community-level Observations. *Public Health Nurs*;33(3):256-63.
- Topaz e.a. (2014). The Omaha System: a systematic review of the recent literature. *J Am Med Inform Assoc*;21(1):163-70.

PACSLAC

- Chan e.a. (2014). Evidence-based development and initial validation of the pain assessment checklist for seniors with limited ability to communicate-II (PACSLAC-II). *Clin J Pain*;30(9):816-24.
- Ellis-Smith e.a. (2016). Measures to assess commonly experienced symptoms for people with dementia in long-term care settings: a systematic review. *BMC Med*;26;14:38.
- Liu e.a. (2010). The psychometric qualities of four observational pain tools (OPTs) for the assessment of pain in elderly people with osteoarthritic pain. *J Pain Symptom Manage*;40(4):582-98.
- Ruest e.a. (2017). Can We Quickly and Thoroughly Assess Pain with the PACSLAC-II? A Convergent Validity Study in Long-Term Care Residents Suffering from Dementia. *Pain Manag Nurs*;pii: S1524-9042(17)30388-0. Full text niet beschikbaar.
- Qi e.a. (2012). The psychometric properties, feasibility and utility of behavioural-observation methods in pain assessment of cognitively impaired elderly people in acute and long-term care: A systematic review. *JBI Libr Syst Rev*. 2012;10(17):977-1085. Full text niet beschikbaar.
- Zwakhalen e.a. (2006). The psychometric quality and clinical usefulness of three pain assessment tools for elderly people with dementia. *Pain*;15;126(1-3):210-20.
- Zwakhalen e.a. (2006). Pain in elderly people with severe dementia: a systematic review of behavioural pain assessment tools. *BMC Geriatr*. 2006 Jan 27;6:3.

PAINAD

- Ellis-Smith e.a. (2016). Measures to assess commonly experienced symptoms for people with dementia in long-term care settings: a systematic review. *BMC Med*;26;14:38.
- Liu e.a. (2010). The psychometric qualities of four observational pain tools (OPTs) for the assessment of pain in elderly people with osteoarthritic pain. *J Pain Symptom Manage*;40(4):582-98.
- Mosele e.a. (2012). Psychometric properties of the pain assessment in advanced dementia scale compared to self assessment of pain in elderly patients. *Dement Geriatr Cogn Disord*;34(1):38-43. Full text niet beschikbaar.
- Qi e.a. (2012). The psychometric properties, feasibility and utility of behavioural-observation methods in pain assessment of cognitively impaired elderly people in acute and long-term care: A systematic review. *JBI Libr Syst Rev*. 2012;10(17):977-1085. Full text niet beschikbaar.
- Warden e.a. (2003). Development and psychometric evaluation of the Pain Assessment in Advanced Dementia (PAINAD) scale. *J Am Med Dir Assoc*;4(1):9-15.
- Zwakhalen e.a. (2006). The psychometric quality and clinical usefulness of three pain assessment tools for elderly people with dementia. *Pain*;15;126(1-3):210-20.
- Zwakhalen e.a. (2006). Pain in elderly people with severe dementia: a systematic review of behavioural pain assessment tools. *BMC Geriatr*. 2006 Jan 27;6:3.

Pain disability index

- Chapman (2011). Evaluating common outcomes for measuring treatment success for chronic low back pain. *Spine (Phila Pa 1976)*;36(21 Suppl):S54-68.
- Ramasamy e.a. (2017). Assessment of Patient-Reported Outcome Instruments to Assess Chronic Low Back Pain. *Pain Med*;18(6):1098-1110.
- Soer e.a. (2013). Extensive validation of the pain disability index in 3 groups of patients with musculoskeletal pain. *Spine (Phila Pa 1976)*;38(9):E562-8.

Pallialine

- Geraadpleegd mei 2017 via: <http://www.pallialine.nl/>. Deze website bevat "richtlijnen palliatieve zorg volgens recente inzichten in diagnostiek en behandeling bij palliatieve zorg beschreven. De richtlijnen zijn multidisciplinair en waar mogelijk evidence based opgesteld".

PalliArts

- Praktische app met landelijke en regionale informatie. Gebaseerd op richtlijnen palliatieve zorg.

PES(DIE)

- Gordon (1994). Nursing Diagnosis: Process and Application, 3ed. Mosby, St Louis.
- Geraadpleegd juni 2017 via: <http://kb.nanda.org/article/AA-00269/0/Our-hospital-requires-us-to-use-the-PES-format-scheme.-How-was-PES-developed-and-what-are-its-origins.html>: "PES" is an acronym that stands for Problem, Etiology (related factors) and Signs/Symptoms (defining characteristics). The PES format was first published by Dr. Marjory Gordon, a founder and former President of NANDA-I. The component parts of NANDA-I diagnoses are now referred to as related factors and defining characteristics, and therefore the wording "PES format" is not used in current NANDA-I books. However, it is still used in several countries and in many publications.

- Müller-Staub e.a. (2007). Meeting the criteria of a nursing diagnosis classification: Evaluation of ICNP, ICF, NANDA and ZEFP. *Int J Nurs Stud*;44(5):702-13.
- Müller-Staub e.a. (2007). Nursing diagnoses, interventions and outcomes--application and impact on nursing practice: a systematic literature review. *Pflege*;20(6):352-71. *Volledige tekst van artikel niet online te vinden.*

Pijn coping inventarisatie schaal

- Kraaimaat e.a. (1997). Pijn-coping-strategieën bij chronische pijnpatiënten: de ontwikkeling van de Pijn-Coping-Inventarisatielijst (PCI). *Gedragstherapie*;30:185-201.
- Kraaimaat & Evers (2003). Pain-coping strategies in chronic pain patients: psychometric characteristics of the pain-coping inventory (PCI). *Int J Behav Med*. 2003;10(4):343-63.
- Perrot e.a. (2008). Active or passive pain coping strategies in hip and knee osteoarthritis? Results of a national survey of 4,719 patients in a primary care setting. *Arthritis Rheum*;15;59(11):1555-62.

PRAFAB Protection, Amount, Frequency, Adjustment en Body Image

- Mulders e.a. (1990). De Inco-test. *Medicus*. (original article; article not found online)
- Hendriks e.a. (2008). Factorial validity and internal consistency of the PRAFAB questionnaire in women with stress urinary incontinence. *BMC Urol*;24:8:1.
- Hendriks e.a. (2007), The psychometric properties of the PRAFAB-questionnaire: a brief assessment questionnaire to evaluate severity of urinary incontinence in women. *BMC*;26:998-1007
- Hendriks e.a. (2008). The minimal important change of the PRAFAB questionnaire in women with stress urinary incontinence: results from a prospective cohort study. *Neurourol Urodyn*;27(5):379-87.

PRISMA-7

- Apóstolo e.a. (2017). Predicting risk and outcomes for frail older adults: an umbrella review of frailty screening tools. *JBJS Database System Rev Implement Rep*;15(4):1154-1208.
- Clegg e.a. (2015). Diagnostic test accuracy of simple instruments for identifying frailty in community-dwelling older people: a systematic review. *Age Ageing*;44(1):148-52.
- Dent e.a. (2016). Frailty measurement in research and clinical practice: A review. *Eur J Intern Med*;31:3-10.
- Hoogendijk e.a. (2013). The identification of frail older adults in primary care: comparing the accuracy of five simple instruments. *Age Ageing*;42(2):262-5.
- Raïche e.a. (2008). PRISMA-7: A case-finding tool to identify older adults with moderate to severe disabilities. *Arch Gerontol Geriatr*;47(1):9-18.
- Sutorius e.a. (2016). Comparison of 10 single and stepped methods to identify frail older persons in primary care: diagnostic and prognostic accuracy. *BMC Fam Pract*;3;17:102.

Protocol Gebruikelijke Zorg Beuningen

- Protocol Gebruikelijke Zorg Gemeente Beuningen (2015). Geraadpleegd mei 2017 via:
decentrale.regelgeving.overheid.nl/cvdr/images/Beuningen/i251510.pdf

Protocol Gebruikelijke Zorg

- Protocol Gebruikelijke zorg (2005). CIZ Centrum Indicatiestelling Zorg.

RAlview

- Achterberg e.a. (2001). Improvements in the quality of co-ordination of nursing care following implementation of the Resident Assessment Instrument in Dutch nursing homes. *J Adv Nurs*;35:268-75.
- Achterberg e.a. (1999). Effects of the Resident Assessment Instrument on the care process and health outcomes in nursing home. A review of the literature. *Scand J Rehabil Med*;31:131-137.
- Berg e.a. (2009). and the ICF: opportunity for operationalizing the ICF. *BMC Health Serv Res*;7:9:47.
- Roth e.a. (2014). The long and bumpy road to outcome-oriented management of long-term care in Germany: implementation of the Resident Assessment Instrument in home-care services. *Int J Health Plann Manage*;29(3):316-29.
- Stolle e.a. (2015). Improving health status and reduction of institutionalization in long-term care--Effects of the Resident Assessment Instrument-Home Care by degree of implementation. *Int J Nurs Pract*;21(5):612-21.
- Stolle e.a. (2012). Effects of the Resident Assessment Instrument in home care settings: results of a cluster randomized controlled trial. *Z Gerontol Geriatr*;45(4):315-22.

RAND-36/SF-36

- Aaronson e.a. (1998). Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J Clin Epidemiol*;51(11):1055-68.
- Hays & Morales (2001). The RAND-36 measure of health-related quality of life. *Ann Med*;33(5):350-7.
- Van der Zee & Sanderman (2012). Het meten van de algemene gezondheidstoestand met de Rand-36. Een handleiding. Research Institute SHARE, UMCG / Rijksuniversiteit Groningen, 2^e druk.

RASS Richmond Agitation and Sedation Scale

- Robinson e.a. (2013). Psychometric analysis of subjective sedation scales in critically ill adults. *Crit Care Med*. 2013 Sep;41(9 Suppl 1):S16-29.
- Varndell e.a. (2015). The validity, reliability, responsiveness and applicability of observation sedation-scoring instruments for use with adult patients in the emergency department: a systematic literature review. *Australas Emerg Nurs J*;18(1):1-23.

REPOS Rotterdam Elderly Pain Observation Scale

- van Herk e.a. (2007). Observation scales for pain assessment in older adults with cognitive impairments or communication difficulties. *Nurs Res*. 2007 Jan-Feb;56(1):34-43. Full text niet beschikbaar.

Richtlijn verslaglegging

- Richtlijn Verpleegkundige en verzorgende verslaglegging (2011). Verpleegkundigen & Verzorgenden Nederland (V&VN), Utrecht.
- Basis voor de inhoud van de richtlijn zijn literatuuronderzoek en input van expertgroepleden.

Risicoscan 2.0 (gezondheidsproblemen)

- Gebaseerd op de Verkorte checklist Veilige Zorg Risicosignalering.

Rode vlaggenlijst voor medicatiegebruik

- Sino (2011). Rode vlaggenlijst voor medicatiegebruik in de thuiszorg. Wat heb jij gezien? Kenniscentrum Innovatie van Zorgverlening Hogeschool Utrecht.
- Sino (2013). Wapperen van vlag legt medicatieprobleem bloot. Tijdschr LVW;2013;12:1:8-9.
- Sino e.a. (2013). Signs and symptoms indicative of potential adverse drug reactions in homecare patients. J Am Med Dir Assoc. 2013;14(12):920-5.
- Sino (2013). Medication management in homecare patients. Dissertation. Hogeschool Utrecht University of Applied Sciences.

Rode Vlaggen App

- Nog in ontwikkeling. In 2016 uitgerold bij twintig wijkteams in Noord-Holland.

SAMPC / SFMPC

- Kinebanian e.a. (1999). Grondslagen van de ergotherapie, Maarssen, Elsevier/ de Tijdstroom Hoofdstuk 10; COPM.
- Geen wetenschappelijke literatuur gevonden.

SBAR Situation, Background, Assessment, Recommendation

- Achrekar e.a. (2016). Introduction of Situation, Background, Assessment, Recommendation into Nursing Practice: A Prospective Study. Asia Pac J Oncol Nurs;3(1):45-50.
- Andreoli e.a. (2010). Using SBAR to communicate falls risk and management in inter-professional rehabilitation teams. Healthc Q;2010;13 Spec No:94-101.
- Boaro e.a. (2010). Using SBAR to improve communication in interprofessional rehabilitation teams. J Interprof Care;24(1):111-4.
- Cornell e.a. (2014). Improving situation awareness and patient outcomes through interdisciplinary rounding and structured communication. J Nurs Adm;44(3):164-9.
- Haig e.a. (2006). SBAR: a shared mental model for improving communication between clinicians. Jt Comm J Qual Patient Saf;32(3):167-75. *Volledekte tekst van artikel niet online te vinden.*
- Lee e.a. (2016). SBAR: towards a common interprofessional team-based communication tool. Med Educ;50(11):1167-1168.

Signaleringskaart eenzaamheid herkennen

- Ontwikkeld door: Academisch werkplaats dementie en Van Kleefinstituut. Geraadpleegd op: <https://www.vankleefinstituut.nl/tools/signaleringskaartje-dementie/>

SMAS Self-Management Ability Scale

- Cramm & Nieboer (2017). Self-management abilities and quality of life among frail community-dwelling individuals: the role of community nurses in the Netherlands. Health Soc Care Community;25(2):394-401.
- Cramm e.a. (2014). Self-management abilities and frailty are important for healthy aging among community-dwelling older people; a cross-sectional study. BMC Geriatr;14:28.
- Cramm e.a. (2012). The relationship between older adults' self-management abilities, well-being and depression. Eur J Ageing;9(4):353-360.
- Cramm e.a. (2012). Validation of the self-management ability scale (SMAS) and development and validation of a shorter scale (SMAS-S) among older patients shortly after hospitalisation. Health Qual Life Outcomes;10:9.
- Schuurmans e.a. (2005). How to measure self-management abilities in older people by self-report: the development of the SMAS-30. Qual Life Res;14(10):2215-2228.
- Steverink e.a. (2005). How to understand and improve older people's selfmanagement of wellbeing. Eur J Aging;2(4):235-244.
- Steverink (2009). Self-Management Ability Scale: SMAS-30/versie 2: achtergrond, handleiding en scoring. 2009. Available from: http://www.nardisteverink.nl/materials/SMAS-30-versie202_achtergrond_handleiding_en_scoring_NSteverink_dec09.pdf

SNAQ / SNAQ65+ Short Nutritional Assessment Questionnaire

- Andraeae e.a. (2015). Psychometric Evaluation of Two Appetite Questionnaires in Patients With Heart Failure. J Card Fail.;21(12):954-8
- Anthony (2008). Nutrition screening tools for hospitalized patients. Nutr Clin Pract;23(4):373-82.
- Kruizenga (2010). The SNAQ(RC), an easy traffic light system as a first step in the recognition of undernutrition in residential care. J Nutr Health Aging;14(2):83-9.
- Phillips e.a. (2010). Nutritional screening in community-dwelling older adults: a systematic literature review. Asia Pac J Clin Nutr;19(3):440-9.
- Rolland (2012). Screening older people at risk of malnutrition or malnourished using the Simplified Nutritional Appetite Questionnaire (SNAQ): a comparison with the Mini-Nutritional Assessment (MNA) tool. J Am Med Dir Assoc;13(1):31-4.
- Van Venrooij e.a. (2007). Quick-and-easy nutritional screening tools to detect disease-related undernutrition in hospital in- and outpatient setting: A systematic review of sensitivity and specificity. Eur Ej Clin Nutr Metab;2:21-37.
- Yaxley e.a. (2015). Identifying Malnutrition in an Elderly Ambulatory Rehabilitation Population: Agreement between Mini Nutritional Assessment and Validated Screening Tools. Healthcare (Basel);11;3(3):822-9.

SOFA-model

- In voor Mantelzorg (2016). Achtergrondinformatie SOFA-model.
- Geen wetenschappelijke literatuur gevonden.

Stakeholder mapping

- MDP Amsta, module Stakeholders voor afdelingshoofden en teamleiders, Oefening Stakeholdersanalyse en –mapping (2013). Amsta.
- Brugha & Varvasovszky (2000). Stakeholder analysis: a review. *Health Policy Plan*;15(3):239-46.
- Shirey (2012). Stakeholder analysis and mapping as targeted communication strategy. *J Nurs Adm*;42(9):399-403.

Stappenplan wijkanalyse maken

- Stappenplan wijkanalyse maken. Van Kleefinstituut.
- Geen wetenschappelijke literatuur gevonden.

Tilburg Frailty Indicator (TFI)

- Apóstolo e.a. (2017). Predicting risk and outcomes for frail older adults: an umbrella review of frailty screening tools. *JBIM Database System Rev Implement Rep*;15(4):1154-1208.
- Daniels e.a. (2012). The predictive validity of three self-report screening instruments for identifying frail older people in the community. *BMC Public Health*;23;12:69.
- Dent e.a. (2016). Frailty measurement in research and clinical practice: A review. *Eur J Intern Med*;31:3-10.
- Gobbens e.a. (2010). The Tilburg Frailty Indicator: psychometric properties. *J Am Med Dir Assoc*;11(5): 344-55.
- Gobbens (2014). The prediction of disability by self-reported physical frailty components of the Tilburg Frailty Indicator (TFI). *Arch Gerontol Geriatr*;59(2):280-7.
- Metzelthin e.a. (2010). The psychometric properties of three self-report screening instruments for identifying frail older people in the community. *BMC Public Health*;31;10:176.
- Pialoux (2012). Screening tools for frailty in primary health care: a systematic review. *Geriatr Gerontol Int*;12(2):189-97.
- Sutton e.a. (2016). Psychometric properties of multicomponent tools designed to assess frailty in older adults: A systematic review. *BMC Geriatr*;29;16:55.

Toolbox Professionele Interactie Dementie

- Stegge e.a. (2016). Professionele interactie in zorgnetwerken van mensen met dementie. Een toolbox voor eerstelijnsprofessionals. Zwolle: Hogeschool Windesheim, ProMemo Expertisecentrum Dementie voor Professionals.
- Deels gebaseerd op bestaande instrumenten. Geen wetenschappelijke literatuur gevonden over de Toolbox als geheel.

Tools voor planning en teamcommunicatie

- Geraadpleegd mei 2017 via: <http://www.nedap-healthcare.com/ouderenzorg/ons-voor-het-team/zelf-organiseren alles-mobiel-bij-de-hand>.

TRAZAG

- TRAZAG TRAnsmuraal Zorg Assessment Geriatrie Complexe zorg voor thuiswonende ouderen (2011). Academisch ziekenhuis Maastricht / MUMC+, Vierde druk.
- Warnier e.a. (2007). Complexe zorg voor ouderen, beoordelingsinstrument voor de praktijkverpleegkundige. *Tijdschr praktijkondersteuning*;2:50-53

U-Care toolkit kwetsbare ouderen

- Bleijenberg e.a. (2012). Toolkit kwetsbare ouderen - U-CARE. Screeningsinstrument en evidence-based zorgplannen voor kwetsbare ouderen. Voor praktijkverpleegkundigen ouderenzorg in de huisartspraktijk.

Valrisico intake

- Intake valrisico. Pilot zorgvoorbeter project valpreventie (2007).
- Geen wetenschappelijke literatuur gevonden.

Valanalyse inventarisatie valrisico 65+ door de eerstelijnszorg

- VeiligheidNL en samenwerkingspartners. Te downloaden via: <https://www veiligheid.nl/valpreventie/interventies/screening/valanalyse>.

Valrisico-inventarisatie

- Valrisico-inventarisatie (VRI). Gemiva (2013).
- Kalkman (2011). Verantwoordingsdocument onderzoek naar de Valrisico-inventarisatielijst. Gemiva-SVG groep.

VAS

- Hjermstad e.a. (2011). Studies comparing Numerical Rating Scales, Verbal Rating Scales, and Visual Analogue Scales for assessment of pain intensity in adults: a systematic literature review. *J Pain Symptom Manage*;41(6):1073-93.
- Schoch e.a. (2017). Patient-Reported Outcome Measures in Pruritus: A Systematic Review of Measurement Properties. *J Invest Dermatol*;pii: S0022-202X(17)31605-6.

Veilige principes in de medicatieketen

- Veilige principes in de medicatieketen (Versie 2, november 2016). Ontwikkeld in het kader van het project 'Veilige principes verder in praktijk', mede mogelijk gemaakt door ZonMw, programma Goed Gebruik Geneesmiddelen. © 2012, 2016 ActiZ, KNMP, NVZA, Verenso, BTN, LHV, LOC, NHG, Patiëntenfederatie Nederland, V&VN.
- Geen wetenschappelijke literatuur gevonden.

Verkorte checklist Veilige Zorg Risicosignalering

- Verkort checklist Veilige Zorg (2015). Zorg voor Beter.
- Gebaseerd op:
 - Richtlijn Screening en behandeling van ondervoeding (Stuurgroep ondervoeding, 2011)
 - SNAQ-RC (Stuurgroep ondervoeding)
 - BEM (Beheer Eigen Medicatie) van het IVM en aantal medicatieveiligheidslijsten van organisaties uit de zorg
 - Richtlijn Mondzorg voor zorgafhankelijke cliënten in verpleeghuizen (NVVA, KNMT en NVG)
 - Richtlijn urine-incontinentie bij kwetsbare ouderen (V&VN en LEVV, 2010)
 - De eenzaamheidsschaal (De Jong Gierveld en Kamphuis, 1985)
 - Screeningschaal DOSS (Bureau voor Toegepaste Sociale Gerontologie, 2001)
 - Richtlijn Herkenning en behandeling van chronische pijn bij kwetsbare ouderen (Verenso, 2011)
 - Folder 'Depressie bij ouderen' (Fonds psychische gezondheid, 2007)
 - Kwaliteitsdocument sector verpleging, verzorging en zorg thuis, indicator 4.4 (VV&T, 2013)
 - GDS: Geriatric Depression Scale (Brink e.a., 1985)
 - Landelijke multidisciplinaire richtlijn Decubitus preventie en behandeling (V&VN, 2011)
 - Folder Oog- en oorproblemen bij ouderen (Het PON, 2010)

Vijf fases van rouwverwerking van Elisabeth Kübler-Ross

- http://www.kublerrossvlaanderen.be/?item=1_de_vijf_fasen_van_rouw
- Geen wetenschappelijke literatuur gevonden.

Vragenlijst Dagelijks Functioneren

- Bijlsma e.a. (1990). Development and validation of the Dutch questionnaire capacities of daily life (VDF) for patients with rheumatoid arthritis. Journal onbekend en full text niet beschikbaar.
- Castrejón e.a. (2011). Clinical composite measures of disease activity for diagnosis and followup of undifferentiated peripheral inflammatory arthritis: a systematic review. J Rheumatol Suppl;87:48-53.
- Kwan (2016). Validity and reliability of the Health Assessment Questionnaire among patients with spondyloarthritis in Singapore. Int J Rheum Dis. [Epub ahead of print]
- Norton e.a. (2014). Health Assessment Questionnaire disability progression in early rheumatoid arthritis: systematic review and analysis of two inception cohorts. Semin Arthritis Rheum;44(2):131-44.
- Oude Voshaar (2011). Measurement properties of physical function scales validated for use in patients with rheumatoid arthritis: a systematic review of the literature. Health Qual Life Outcomes;7:9:99.

Vragenlijst Rouw (in: Richtlijn Rouw, IKNL)

- Boelen e.a. (2017). The Traumatic Grief Inventory Self-Report Version (TGI-SR): Introduction and Preliminary Psychometric Properties. J Loss Trauma;22:196:212.
- Boelen e.a. (2003). Reliability and validity of the Dutch version of the inventory of traumatic grief (ITG). Death Stud;27(3):227-47.
- Boelen e.a. (2001). Psychometrische eigenschappen van de Rouw VragenLijst (RVL). Gedrag & Gezondheid;29:172-185.

Vroegsignalering Dementie

- Veenstra (2010). Vroegsignalering helpt cliënt én mantelzorger. Academische werkplaats Dementie. Tijdschrift LVW;10;3;12-14.
- Geen wetenschappelijke literatuur gevonden.

Waaijer medicatieveiligheid

- Medicatiewaaijer: Medicatiezorg, praktische informatie en tips (2013). Vilans, geraadpleegd april 2017 via: www.zorgvoorbeter.nl
- Geen wetenschappelijke literatuur gevonden.

Wondzorg app

- Boomerweb. Te downloaden via: <https://play.google.com/store/apps/details?id=nl.boomerweb.wondzorgapp>.

Zarit-12

- Al-Rawashdeh e.a. (2016). Psychometrics of the Zarit Burden Interview in Caregivers of Patients With Heart Failure. J Cardiovasc Nurs;31(6):E21-E28.
- Hagell e.a. (2017). Assessment of Burden Among Family Caregivers of People With Parkinson's Disease Using the Zarit Burden Interview. J Pain Symptom Manage;53(2):272-278
- Higginson e.a. (2010). Short-form Zarit Caregiver Burden Interviews were valid in advanced conditions. J Clin Epidemiol;63(5):535-42.
- Lin (2017). Measuring burden in dementia caregivers: Confirmatory factor analysis for short forms of the Zarit Burden Interview. Arch Gerontol Geriatr;68:8-13.
- Van Durme e.a. (2012). Tools for measuring the impact of informal caregiving of the elderly: a literature review. Int J Nurs Stud;49(4):490-504.
- Whalen & Buchholz. (2009). The reliability, validity and feasibility of tools used to screen for caregiver burden: a systematic review. JBI Libr Syst Rev;7(32):1373-1430.

Zelfmanagement ondersteuning chronisch zieken

- Ondersteun zelfmanagement van chronisch zieken (2011). Vilans.
- In rapport verwijzing naar meerdere referenties.

Zelfredzaamheidsmatrix

- Bannink e.a. (2015). Psychometric properties of self-sufficiency assessment tools in adolescents in vocational education. *BMC Psychol*;25;3:33.
- Fassaert e.a. (2014). Psychometric properties of the Dutch version of the self-sufficiency matrix (SSM-D). *Community Ment Health J*;50(5):583-90.
- Mast e.a. (2014). Zo zelfredzaam. Een overzicht van instrumenten voor het meten van zelfredzaamheid. Vilans.
- Van den Boom e.a. (2016). Psychometrische kenmerken van de Zelfredzaamheid-Matrix (ZRM) en het ZRM-Supplement: Ouderschap. *Kind Adolesc*;37:105–119.

Zelfredzaamheidsmeter

- Mast e.a. (2014). Zo zelfredzaam. Een overzicht van instrumenten voor het meten van zelfredzaamheid. Vilans.
- Zelfredzaamheid | Zelfredzaamheidsmeter | © Vilans, juni 2013.
- Zelfredzaamheidsmeter; uitleg. Geraadpleegd mei 2017: <https://www.btsq.nl/infobulletin/zelfredzaamheidsmeter.html>.
- Geen wetenschappelijke literatuur gevonden.

Zelfredzaamheidsmonitor

- Mast e.a. (2014). Zo zelfredzaam. Een overzicht van instrumenten voor het meten van zelfredzaamheid. Vilans.
- Zelfredzaamheidsmonitor (2014). Vilans & CTOW.
- Geen wetenschappelijke literatuur gevonden.

Zelfredzaamheidsradar

- De ZelfredzaamheidsRadar© is gebaseerd op de Care Dependency Scale van Dijkstra et al (1996). Geen wetenschappelijke literatuur gevonden over de ZelfredzaamheidsRadar© zelf.
- Dijkstra e.a. (1996). Nursing-care dependency. Development of an assessment scale for demented and mentally handicapped patients. *Scand J Caring Sci*;0(3):137-43.
- Kottner e.a. (2010). Interrater reliability and agreement of the Care Dependency Scale in the home care setting in the Netherlands. *Scand J Caring Sci*;24 Suppl 1:56-61.
- Mast e.a. (2014). Zo zelfredzaam. Een overzicht van instrumenten voor het meten van zelfredzaamheid. Vilans.

Zelfstandigheidslijst

- Mast e.a. (2014). Zo zelfredzaam. Een overzicht van instrumenten voor het meten van zelfredzaamheid. Vilans.
- Geen wetenschappelijke literatuur gevonden.

Zorgmodule zelfmanagement 1.0

- Zorgmodule Zelfmanagement 1.0. Het ondersteunen van eigen regie bij mensen met één of meerdere chronische ziekten (2014). CBO.
- De zorgmodule is gebaseerd op praktijkervaringen, aangevuld met beschikbare wetenschappelijke evidentie, en consensus in de ontwikkelgroep.

Zorgpad Stervensfase

- IKNL, geraadpleegd mei 2017 via <https://www.iknl.nl/palliatieve-zorg/verbetertrajecten/zorgpad-stervensfase>. “Het Zorgpad Stervensfase is onderdeel van de richtlijn Zorg in de stervensfase. Het zorgpad is gericht op maximaal comfort voor patiënten en naasten tijdens de stervensfase.”

Zorgstandaard Dementie

- Zorgstandaard Dementie, Steun en Zorg (2013). Alzheimer Nederland & Vilans. Voor de totstandkoming van De Zorgstandaard Dementie is gebruik gemaakt van recente wetenschappelijke inzichten en geldende richtlijnen van verschillende professionele beroepsgroepen, zie bronnenlijst in document.

3 manieren om stakeholders te identificeren

- 3 Eenvoudige manieren om uw stakeholders te identificeren. In voor Zorg, 2016. Gebaseerd op de bronnen: Hooge, E, F.J. de Vijlder en M. van der Sluis (2005) “Stakeholders in beeld”. Amsterdam, Max Goote Kenniscentrum en Kennisnet: Instrumenten voor het identificeren van en communiceren met stakeholders.